

Table 12. Yield summary of the irrigated cotton variety test at the Texas Agricultural Experiment Station, Lubbock, TX, 2002.

Designation	1998	1999	2000	2001	2002	Comp.	
						Average	Average*
Five Year Average							
All-Tex Atlas	833	737	676	839	922	801	
Paymaster 2200 RR	929	821	693	916	939	860	
Paymaster 2280 BG/RR	621	773	742	1072	895	821	
Paymaster 2326 RR	879	790	693	890	901	831	
Paymaster 2326 BG/RR	819	786	755	946	872	836	
Tamcot Luxor	736	782	720	858	628	745	
Tamcot Sphinx	695	892	830	860	630	781	
Four Year Average							
FiberMax 819		823	874	847	756	825	819
Three Year Average							
FiberMax 958			861	960	1035	952	940
FiberMax 5013			715	794	833	781	769
FiberMax 5015			790	909	1000	900	888
FiberMax 5017			667	893	826	795	784
Paymaster 2379 RR			790	888	906	861	850
Phytogen 355			812	1065	787	888	876
Stoneville 2454 R			866	866	770	834	822
Stoneville 3539 BR			777	1028	852	886	874
Stoneville 4793 R			799	1017	795	870	859
Stoneville 4892 BR			905	1161	868	978	966
Stoneville 5599 BR			989	1192	884	1022	1010

\*Patterson, R.E. 1950. A method of adjustment for calculating comparable yields in variety tests.