

ACKNOWLEDGMENTS

Fiber Properties were determined at the International Textile Center, Texas Tech University with financial support from the Texas Food and Fiber Commission and the Plains Cotton Improvement Program.

The financial and moral support provided by the Plains Cotton Improvement Program is invaluable as we strive to become a consistent supplier of high quality cotton.

Plot harvesting, sample ginning, and data collection were performed by: Jerry Baeza, Bailey Boyd, Amanda Burns, Trey Carter, Billy Chaffin, Leigh Cranmer, Dusty Creel, Michael Drachenberg, Erik Everitt, Tom Fouts, Johnny Fuentes, Anna Hall, Casey Hardin, Lisa Harrell, Ryan Heinrich, Bryan Henson, Shauna Lewis, Kyle Long, Carol Mason, Valerie Morgan, Meredith Pharr, Jody Rowland, Lyndon Schoenhals, Monica Sheehan, Ross Stegemoeller, Clay Teeter, Raymond Tillis, Jason Trammell, Will Trammell, Whitney Verett, Paul Ward, Leslie Wells, Jennifer White, Jerry White and Steven White. Sheryl Angerer provided secretarial assistance with the typing and editing of the numerous tables associated with this report. Valerie Morgan and Lyndon Schoenhals performed table preparation and statistical analyses. The assistance of all of these people is gratefully acknowledged and truly appreciated.