

Table 41. Production information for irrigated performance test at the AG-CARES Farm, Lamesa, TX, 2004.

---

Test:	Nematode Variety and Strains				
Planting Date:	May 6				
Row Spacing:	40-in				
Planting Pattern:	Solid in Rye Stubble				
Herbicide:	Prowl @ 3.0 pt/A applied on April 22 CornerStone @ 1 qt/A applied on May 24				
Fertilizer:	32-0-0 @ 50 lbs/A applied on June 9 32-0-0 @ 25 lbs/A applied on July 2 and 22				
Irrigations: (center pivot)	1.5 inches pre-plant				
	May	June	July	August	September
	7 - .3 in.	4 - .5 in.	2 - .6 in.	2 - .5 in.	1 - .5 in
	9 - .3 in.	9 - .75 in	5 - .6 in.	23 - .5 in	
	10 - .3 in.	13 - .55 in	8 - .5 in.	27 - .5 in.	
	12 - .3 in.	16 - .55 in	12 - .5 in		
	14 - .75 in.	24 - .44 in	15 - .5 in.		
	16 - .3 in.	27 - .55 in	19 - .5 in.		
	19 - .3 in		22 - .5 in.		
Insecticide:	Temik @ 4.0 lbs/A at planting				
Harvest Aids:	Boll'd @ 16 oz/A + Ginstar @ 4.0 oz/A on September 27 Cyclone @ 24 oz/A on October 9				
Harvest Date:	November 9				
Freeze Date:	November 30				