

Table 53. Production information for irrigated performance test at the AG-CARES Farm, Lamesa, TX, 2005.

Test:	Nematode Variety and Strains				
Planting Date:	May 9				
Row Spacing:	40-in				
Planting Pattern:	Solid in Rye Stubble				
Herbicide:	Prowl @ 3.0 pt/A applied on April 27 Roundup Weathermax @ 22 oz/A applied on April 30 Roundup @ 1 qt/A on June 6				
Fertilizer:	32-0-0 @ 50 lbs/A applied on June 9 32-0-0 @ 25 lbs/A applied on July 2 and 22				
Irrigations: (center pivot)	1.75 inches pre-plant				
	May	June	July	August	September
	13 - .4 in.	14 - .4 in.	2 - .44 in.	4 - .44 in.	3 - .44 in.
	20 - .4 in.	17 - .88 in.	5 - .44 in.	11 - .44 in.	8 - .44 in.
	24 - .4 in.	20 - .44 in.	8 - .44 in.	14 - .44 in.	
		23 - .55 in.	10 - .44 in.	23 - .44 in.	
		26 - .44 in.	19 - .44 in.	26 - .44 in.	
		29 - .44 in.	23 - .44 in.	29 - .44 in.	
			26 - .44 in.		
Insecticide:	Temik @ 3.5 lbs/A at planting				
Harvest Aids:	Prep @ 1 pt/A + Dep @ 1 pt/A on September 27 Gramoxone Max @ 16 oz/A on October 12				
Harvest Date:	October 18				
Freeze Date:	November 15				